

# A Note on Stable Quadratic Polynomials over Fields of Characteristic Two

OMRAN AHMADI

Claude Shannon Institute

University College Dublin

Dublin 4, Ireland

omran.ahmadi@ucd.ie

October 23, 2009

## Abstract

In this note, first we show that there is no stable quadratic polynomial over finite fields of characteristic two and then show that there exist stable quadratic polynomials over function fields of characteristic two.

## 1 Introduction

Let  $K$  be a field and  $K[x]$  be the polynomial ring over  $K$ . A polynomial  $f(x) \in K[x]$  is called stable if  $f(x), ff(x), fff(x), \dots, f^{(n)}(x), \dots$  is a sequence of irreducible polynomials in  $K[x]$ . Recently there has been an interest in the study of stable polynomials (see [1, 2, 3, 5, 7, 8, 9])

In [9], Ostafe and Shparlinski studied stable quadratic polynomials over finite fields and obtained an upper bound on the length of the critical orbits of stable quadratic polynomials over finite fields with odd characteristic. They also posed the question of estimating the number of stable quadratic polynomials over finite fields. Here we show that there is no stable quadratic polynomial over finite fields of characteristic two by proving the following theorem.

**Theorem 1.** *Let  $q = 2^m$ , and let  $f(x) = cx^2 + ax + b \in \mathbb{F}_q[x]$  where  $\mathbb{F}_q$  denotes the field with  $q$  elements and  $a, b \in \mathbb{F}_q^*$ . Then  $fff(x)$  cannot be an irreducible polynomial over  $\mathbb{F}_q$ .*

## 2 Proof of Theorem 1

For simplicity we assume that  $c = 1$  and hence  $f(x)$  is a monic polynomial. The same proof can be used for the non-monic polynomials. We also note that in what follows  $\text{Tr}_{|K||L|}(\cdot)$  denotes the trace map from field  $K$  with  $|K|$  elements to its subfield  $L$  with  $|L|$  elements. Our proof is based on the following two well-known lemmas.

**Lemma 2.** *[4, Corollary 3.6] Let  $q = 2^m$ , and let  $f(x) = x^2 + ax + b \in \mathbb{F}_q[x]$  where  $\mathbb{F}_q$  denotes the field with  $q$  elements and  $a, b \in \mathbb{F}_q^*$ . Then  $f(x)$  is irreducible over  $\mathbb{F}_q$  if and only if  $\text{Tr}_{q|2}(\frac{b}{a^2}) = 1$ .*

The following lemma is known as Capelli's lemma and can be found in [6] too.

**Lemma 3.** *Let  $f(x)$  be a degree  $n$  irreducible polynomial over  $\mathbb{F}_q$ , and let  $g(x) \in \mathbb{F}_q[x]$ . Then  $p(x) = f(g(x))$  is irreducible over  $\mathbb{F}_q$  if and only if for some root  $\alpha$  of  $f(x)$  in  $\mathbb{F}_{q^n}$ ,  $g(x) - \alpha$  is an irreducible polynomial over  $\mathbb{F}_{q^n}$ .*

Now suppose that both  $f(x) = x^2 + ax + b$  and  $ff(x)$  are irreducible over  $\mathbb{F}_q$ . We show that  $fff(x)$  cannot be an irreducible polynomial over  $\mathbb{F}_q$ . Suppose for the contrary that  $fff(x)$  is irreducible over  $\mathbb{F}_q$ . Using Lemma 3,  $fff(x)$  is irreducible over  $\mathbb{F}_q$  if and only if  $h(x) = f(x) - \alpha$  is irreducible over  $\mathbb{F}_{q^4}$  for some root  $\alpha$  of  $ff(x)$  in  $\mathbb{F}_{q^4}$ . Applying Lemma 2,  $h(x)$  is irreducible over  $\mathbb{F}_{q^4}$  if and only if  $\text{Tr}_{q^4|2}(\frac{b-\alpha}{a^2}) = 1$ . Using the properties of the trace map in one hand

$$\text{Tr}_{q^4|2}\left(\frac{b-\alpha}{a^2}\right) = \text{Tr}_{q^4|2}\left(\frac{b}{a^2}\right) - \text{Tr}_{q^4|2}\left(\frac{\alpha}{a^2}\right) \quad (1)$$

and on the other hand

$$\text{Tr}_{q^4|2}\left(\frac{b}{a^2}\right) = 4\text{Tr}_{q|2}\left(\frac{b}{a^2}\right) = 0.$$

Thus from above and Equation 1, we conclude that  $\text{Tr}_{q^4|2} \left( \frac{\alpha}{a^2} \right) = 1$ . But

$$\begin{aligned} \text{Tr}_{q^4|2} \left( \frac{\alpha}{a^2} \right) &= \text{Tr}_{q|2} \left( \text{Tr}_{q^4|q} \left( \frac{\alpha}{a^2} \right) \right) \\ &= \text{Tr}_{q|2} \left( \frac{\text{Tr}_{q^4|q}(\alpha)}{a^2} \right). \end{aligned}$$

Now suppose that  $ff(x) = x^4 + c_3x^3 + c_2x^2 + c_1x + c_0$ . Then it is easy to see that  $c_3 = 0$ . Finally, since  $\alpha$  is a root of  $ff(x)$  in  $\mathbb{F}_{q^4}$ , we deduce that  $\text{Tr}_{q^4|q}(\alpha) = c_3 = 0$ , and hence  $\text{Tr}_{q^4|2} \left( \frac{\alpha}{a^2} \right) = 0$  which is a contradiction.

### 3 The case of function fields of characteristic two

Let  $\mathbb{F}_2(t)$  be the rational function field in  $t$  over  $\mathbb{F}_2$ , where  $t$  is transcendental over  $\mathbb{F}_2$ , and let  $\mathbb{F}_2(t)[x]$  be the polynomial ring over  $\mathbb{F}_2(t)$  in the variable  $x$ .

One expects that a theorem similar to Theorem 1 to hold for quadratic polynomials in  $\mathbb{F}_2(t)[x]$  but this is not true as the following example shows.

Suppose  $f(x) = x^2 + t \in \mathbb{F}_2(t)[x]$ . Then it is easy to see that

$$f^{(n)}(x) = x^{2^n} + t^{2^{n-1}} + t^{2^{n-2}} + \cdots + t^2 + t.$$

Now from Eisenstein's criterion for function fields [10, Proposition III.1.14] it is easy to see that for every  $n \geq 1$ ,  $f^{(n)}(x)$  is irreducible over  $\mathbb{F}_2(t)$  and hence  $f(x)$  is a stable polynomial in  $\mathbb{F}_2(t)[x]$ .

### 4 Acknowledgements

The author would like to thank Igor Shparlinski for helpful comments on an earlier version of this note.

### References

- [1] Nidal Ali. Stabilité des polynômes. *Acta Arith.*, 119(1):53–63, 2005.
- [2] Mohamed Ayad and Donald L. McQuillan. Irreducibility of the iterates of a quadratic polynomial over a field. *Acta Arith.*, 93(1):87–97, 2000.

- [3] Mohamed Ayad and Donald L. McQuillan. Corrections to: “Irreducibility of the iterates of a quadratic polynomial over a field” [Acta Arith. **93** (2000), no. 1, 87–97; MR1760091 (2001c:11031)]. *Acta Arith.*, 99(1):97, 2001.
- [4] Ian. F. Blake, XuHong Gao, Alfred J. Menezes, Ronald C. Mullin, Scott A. Vanstone, and Tomik Yaghoobian. *Application of finite fields*. Kluwer, 1993.
- [5] Nigel Boston and Rafe Jones. Settled polynomials over finite fields. *preprint*, 2009.
- [6] Stephen D. Cohen. On irreducible polynomials of certain types in finite fields. *Proc. Cambridge Philos. Soc.*, 66:335–344, 1969.
- [7] Rafe Jones. Iterated Galois towers, their associated martingales, and the  $p$ -adic Mandelbrot set. *Compos. Math.*, 143(5):1108–1126, 2007.
- [8] Rafe Jones. The density of prime divisors in the arithmetic dynamics of quadratic polynomials. *J. Lond. Math. Soc. (2)*, 78(2):523–544, 2008.
- [9] Alina Ostafe and Igor Shparlinski. On the length of critical orbits of stable quadratic polynomials. *preprint*, 2009.
- [10] Henning Stichtenoth. *Algebraic function fields and codes*. Universitext. Springer-Verlag, Berlin, 1993.